

Survey of Neutrophil to Lymphocyte Ratio as Prognostic Factor in Colorectal Cancer

Negin Farshchian^{*,***}, Maral Soleimani^{**}, Sahel Heydarheydari^{***},
Nazanin Farshchian^{****,**}

^{*}Department of Radiation Oncology, Faculty of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

^{**}Clinical Research Development Center, Imam Reza Hospital, Kermanshah, Iran

^{***}Department of Medical Physics, Faculty of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

^{****}Department of Radiology, Faculty of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Abstract

Background: Neutrophil-lymphocyte ratio (NLR) is derived from the neutrophil and lymphocyte count obtained from routine complete blood count with the differential test (CBC/diff). The aim of the present study is to determine the relationship between NLR with prognostic factors in colorectal cancer.

Material and Methods: A total of 70 patients with colorectal cancer were studied according to the research objectives. Independent T-test and Mann-Whitney U Mann-Whitney test were used to compare the NLR with qualitative variables such as pathology type. Pearson or Spearman correlation coefficient was used for correlating the quantitative variables. The significance level was considered as $P < 0.05$.

Results: Most of the enrolled patients were in the age group of 60-69 years (31.5%). The mean and standard deviation of NLR was 2.1 ± 0.1 . Only 4.3% of the subjects had NLR greater than 5. The association between NLR with N-stage ($P=0.005$) and inflammatory factor (CRP) was statistically significant ($P = 0.016$). The correlation between NLR and these variables showed that Grade ($P=0.091$), PNI ($P=0.093$), LDH ($P=0.20$), diabetes ($P=0.264$), M-stage ($P=0.282$), ESR ($P=0.386$), LSVI ($P=0.473$), T-stage ($P=0.527$), CEA ($P=0.582$), and BMI ($P=0.681$) with NLR was not meaningful.

Conclusion: NLR can be associated with prognosis and risk of disease progression, especially in lymph nodes. Since NLR does not require additional testing and request and is quite easy and cost-effective, it is recommended considering this ratio during the initial assessment and the classification of the risk and stage of the disease.

Keywords: Colorectal cancer (CRC), Neutrophil-lymphocyte ratio (NLR), Prognostic Factor

♦Corresponding Author:

Negin Farshchian, MD
Department of Radiation
Oncology, Kermanshah
University of Medical Sciences,
Clinical Research Development
Center, Imam Reza Hospital,
Kermanshah, Iran
Tel.: +98 9188556393
Email: Farshchian.n@kums.ac.ir

Introduction

Colorectal cancer (CRC) is the fourth most frequent malignancy in men and the third among women.¹ Identification of prognostic factors is an important step to assess clinical outcome in the treatment of CRC. Tumor stage, tumor grade, number of resected lymph nodes, peri-neural or vascular invasion, and also the carcinoembryonic antigen (CEA) level have been introduced as prognostic markers in previous studies.² The circulating neutrophil-lymphocyte ratio (NLR) which is derived from the neutrophil and lymphocyte count, which is obtained from routine complete blood count with the differential test (CBC w/diff). NLR as a reliable and readily available prognostic marker in various tumor types including CRC, reflects a systemic inflammatory response.³ Based on multiple studies, high NLR is associated with low survival in multiple cancers including breast, pancreatic, gastric, renal, and CRC.⁴ Increased NLR has been associated with adverse clinicopathological parameters and poor survival rates in CRC⁵ and suggested as a simple index of systemic inflammatory response in seriously ill patients.⁶

The aim of the present study is to determine the relationship between NLR with prognostic factors in CRC.

Material and Methods

In this descriptive cross-sectional study, 70 patients with CRC were recruited according to the research objectives. A questionnaire was designed for data collection related to the objectives of this study. Information was collected by filling out the forms from patients' files for each individual and some of the variables were obtained with appropriate tests. After completing the data collection form, the information obtained was entered into SPSS software version 20 and analyzed. Descriptive statistics including calculating numerical indices such as mean and standard deviation for quantitative traits and percentage for qualitative traits were used to summarize. Independent T-test and Mann-Whitney U test were used to compare the NLR with

qualitative variables such as pathology type. Pearson or Spearman correlation coefficient was used for correlating the quantitative variables. The significance level was considered as $P < 0.05$. The ethics committee of Kermanshah University of Medical Sciences (KUMS) approved this study (Approved research plan No.: 95016).

Results

According to table 1, most of the patients were in the age group of 60-69 years (31.5%). Also, 47.1% of the enrolled patients were female and 52.9% were male. The mean and standard deviation (SD) of NLR was 2.1 ± 0.1 . Only 4.3% of the subjects had NLR greater than 5. In the age group under 50 years of age, the mean \pm SD of NLR was 9.9 ± 0.8 . The mean \pm SD of NLR was 0.9 ± 0.6 and 2.3 ± 1.1 in the age groups 50-60 and 70-69 years, respectively. Also, in the age group older than 70 years, the mean \pm SD of NLR was 2.4 ± 1.3 . Regarding these results, although with increasing age, the NLR was slightly decreased, the statistical test did not show any significant difference.

As shown in table 1, 50% of the patients had grade 1 tumor. Also, 77.1% had stage 3, 57.1% had no lymph node involvement, and only 4 had metastasis (M1). The correlation between NLR and these variables showed that Grade ($P=0.091$), T-stage ($P=0.527$), and M-stage ($P=0.282$) had no statistically significant relationship with NLR, but the relationship between NLR with N-stage was statistically significant ($P=0.005$). According to table 1, 45.7% of the cases had negative CRP. The mean and standard deviation of ESR was 31.6 ± 24.6 mm/h. The relation between ESR and NLR was not statistically significant ($P=0.386$), but the one between NLR and CRP was statistically significant ($P=0.016$). The association between ESR and CRP was also assessed by NLR.

Results showed that none of the two LSVI factors ($P=0.473$) and PNI ($P=0.093$) had a significant relation with NLR. Table 1 shows that 15.7% of the patients had a mucosal pathology and 84.3% of non-muscular pathology, and only 11.4%

Table 1. Clinical characteristics of the enrolled patients and clinicopathological findings

Gender			LVSI		
	Male	37 (47.1%)		Positive	11 (16%)
	Female	33 (52.9%)		Negative	59 (84%)
Age			PNI		
	<50	13 (18.5%)		Positive	23 (33%)
	50-59	18 (25.7%)		Negative	47 (67%)
	60-69	22 (31.5%)			
	>70	17 (24.2%)			
BMI			LDH	<200	02 (3%)
	<20	05 (7%)		200-300	32 (46%)
	20-25	43 (61.5%)		300-400	28 (40%)
	25-30	19 (27%)		>400	08 (11%)
	>30	03 (4.5%)			
Diabetes			CEA	<3	29 (41.5%)
	Positive	08 (11.5%)		3-6	20 (28.5%)
	Negative	62 (88.5%)		>6	21 (30%)
Type of Pathology			ESR	<20	27
	Mucinous	11 (16%)		(38.5%)	
	Non-mucinous	59 (84%)		20-69	37 (53%)
				>70	06 (8.5%)
Grade			CPR	-	32 (46%)
	1	35 (50%)		+	26 (37%)
	2	29 (41.5%)		++	10 (14%)
	3	06 (8.5%)		+++	02 (3%)
	4	00 (0%)			
T-stage					
	1	01 (1.5%)			
	2	01 (1.5%)			
	3	54 (77%)			
	4	14 (20%)			
Node Involvement					
	Yes	30 (43%)			
	No	40 (47%)			
Metastasis					
	Yes	04 (6%)			
	No	66 (94%)			

of the patients had diabetes. Also, there was no significant relationship between the two pathologic factors ($P=0.827$) and diabetes ($P=0.264$) with NLR. The mean and standard deviation of BMI in the study population was 23.9 ± 3.33 and the highest frequency was in the 25-20 group with 61.4%. The mean and standard deviation of LDH and CEA were 317.75 ± 7.7 and 16.5 ± 39.1 , respectively. The association of these three factors with NLR was investigated and it was found that the relationship between the three, BMI ($P=0.681$), LDH ($P=0.20$), and CEA ($P=0.582$) and NLR was not meaningful.

Discussion

Identifying the prognostic and predictive factors of CRC plays a key role in determining the most effective treatment. However, there are still pending satisfactory detecting prognostic markers. According to study on the prognostic impact of increased NLR in multiple settings of CRC, it was found that increased NLR leads to poor prognosis in CRC and NLR associated with the highest degree of discrimination of outcomes.⁷ Recently, two meta-analyses showed significant evidence to support the association between high

NLR and poor prognosis in patients with CRC.^{8,9} Previous studies have shown the role of albumin, serum CRP, CEA, absolute neutrophil counts, and NLR in predicting the clinical outcomes of patients with CRC diagnosis.¹⁰ Several studies have demonstrated the role of NLR as a prognostic factor in CRC patients.¹¹ Immune cells reflect tumor development while NLR acts closely with the systematic inflammatory response.

In the present study, we surveyed patients with CRC, most of them being stage III and not having metastasis. The relation of stage and CEA with NLR and some variables and a number of other previously expressed variables, were studied as well.

Carruthers et al. found no correlation between NLR and tumor downstaging.¹² However, in another study, it was associated with T4 stage cancer.¹³ Based on the results of the current study, the correlation between NLR and N-stage was statistically significant. Tumor differentiation data were available from three studies involving 1329 CRC patients. Based on the overall estimates, NLR greater than 5 is associated with low tumor differentiation with no meaningful heterogeneity displayed that CRC patients with $NLR > 5$ were expected to be in T4 stage.⁸

Previous studies reported that NLR could be an independent prognostic factor to predict long-term outcomes in stage III and II CRC, and also an important predictive marker in stage IV CRC.² Several researchers have reported the relationship between NLR and CEA level before cancer therapy. The pooled estimates presented a significant relationship between $NLR > 5$ and higher CEA level (> 5). Elsewhere, pooled OR was 1.59 without significant heterogeneity.^{14-16,17} However, according to the data of the present study, the relationship between the three BMI, LDH, and CEA with NLR was not meaningful.

Conclusion

NLR can be associated with prognosis and risk of disease progression, especially in lymph nodes. Since NLR does not require additional

testing and request and is quite easy and cost-effective, it is recommended to consider this ratio during the initial assessment and the classification of the risk and stage of the disease.

The evaluation of the NLR factor can be useful in determining the risk and progression of the disease and should be considered in treatment decisions.

In the current study, due to the limited numbers of patients, the optimal cut-off of NLR was not achieved. It would be possible to achieve an optimal cut-off for this ratio, as a better prognostic marker, in other studies with more sample sizes. Furthermore, it seems necessary to perform similar studies to follow up the survival of patients and determine the relationship between NLR with the survival rate in order to reveal the prognostic importance of this marker.

Acknowledgments

The authors gratefully acknowledge the Research Council of Kermanshah University of Medical Sciences and Clinical Research Development Center, Imam Reza Hospital for their financial support and cooperation. This work (approved research project No: 95016) was performed in partial fulfillment of the requirements for MD degree of Maral Soleimani, in the Faculty of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran.

Conflict of Interest

None declared.

References

1. Madani SH, Sadeghi E, Rezaee A, Sadeghi M, Khazaei S, Amirifard N, et al. Survey of HER2-neu expression in colonic adenocarcinoma in the west of Iran. *Asian Pac J Cancer Prev*. 2015; 16(17): 7671-4.
2. Kim JH, Lee JY, Kim HK, Lee JW, Jung SG, Jung K, et al. Prognostic significance of the neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio in patients with stage III and IV colorectal cancer. *World J Gastroenterol*. 2017; 23(3): 505-15.
3. Absenger G, Szkandera J, Stotz M, Postlmayr U, Pichler M, Röss AL, et al. Preoperative neutrophil-to-lymphocyte ratio predicts clinical outcome in patients

- with stage II and III colon cancer. *Anticancer Res.* 2013; 33(10): 4591-4.
4. Shen L, Zhang H, Liang L, Li G, Fan M, Wu Y, et al. Baseline neutrophil-lymphocyte ratio (≥ 2.8) as a prognostic factor for patients with locally advanced rectal cancer undergoing neoadjuvant chemoradiation. *Radiat Oncol.* 2014;9:295. doi: 10.1186/s13014-014-0295-2.
 5. Guthrie GJ, Roxburgh CS, Richards CH, Horgan PG, McMillan DC. Circulating IL-6 concentrations link tumour necrosis and systemic and local inflammatory responses in patients undergoing resection for colorectal cancer. *Br J Cancer.* 2013; 109(1): 131-7.
 6. Zahorec R. Ratio of neutrophil to lymphocyte counts-Rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl Lek Listy.* 2001; 102: 5-14.
 7. Chen ZY, Raghav K, Lieu CH, Jiang ZQ, Eng C, Vauthey JN, et al. Cytokine profile and prognostic significance of high neutrophil-lymphocyte ratio in colorectal cancer. *Br J Cancer.* 2015; 112(6): 1088-97.
 8. Li MX, Liu XM, Zhang XF, Zhang JF, Wang WL, Zhu Y, et al. Prognostic role of neutrophil-to-lymphocyte ratio in colorectal cancer: a systematic review and meta-analysis. *Int J Cancer.* 2014; 134: 2403-13.
 9. Malietzis G, Giacometti M, Kennedy RH, Athanasiou T, Aziz O, Jenkins JT. The emerging role of neutrophil to lymphocyte ratio in determining colorectal cancer treatment outcomes: a systematic review and meta-analysis. *Ann Surg Oncol.* 2014; 21: 3938-46.
 10. Tsai PL, Su WJ, Leung WH, Lai CT, Liu CK. Neutrophil-lymphocyte ratio and CEA level as prognostic and predictive factors in colorectal cancer: A systematic review and meta-analysis. *J Cancer Res Ther.* 2016; 12(2): 582-9.
 11. Wu Y, Li C, Zhao J, Yang L, Liu F, Zheng H, et al. Neutrophil-to-lymphocyte and platelet-to-lymphocyte ratios predict chemotherapy outcomes and prognosis in patients with colorectal cancer and synchronous liver metastasis. *World J Surg Oncol.* 2016; 14(1): 289.
 12. Carruthers R, Tho LM, Brown J, Kakumanu S, McCartney E, McDonald AC. Systemic inflammatory response is a predictor of outcome in patients undergoing preoperative chemoradiation for locally advanced rectal cancer. *Colorectal Dis.* 2012; 14:e701-7.
 13. Walsh SR, Cook EJ, Goulder F, Justin TA, Keeling NJ. Neutrophil-lymphocyte ratio as a prognostic factor in colorectal cancer. *J Surg Oncol.* 2005; 91: 181-4.
 14. Zhang Y, Peng Z, Chen M, Liu F, Huang J, Xu L, et al. Elevated neutrophil to lymphocyte ratio might predict poor prognosis for colorectal liver metastasis after percutaneous radiofrequency ablation. *Int J Hyperthermia.* 2012; 28: 132-40.
 15. Hung HY, Chen JS, Yeh CY, Changchien CR, Tang R, Hsieh PS, et al. Effect of preoperative neutrophil-lymphocyte ratio on the surgical outcomes of stage II colon cancer patients who do not receive adjuvant chemotherapy. *Int J Colorectal Dis.* 2011; 26: 1059-65.
 16. Kwon HC, Kim SH, Oh SY, Lee S, Lee JH, Choi HJ, et al. Clinical significance of preoperative neutrophil-lymphocyte versus platelet-lymphocyte ratio in patients with operable colorectal cancer. *Biomarkers.* 2012; 17: 216-22.
 17. Halazun KJ, Aldoori A, Malik HZ, Al-Mukhtar A, Prasad KR, Toogood GJ, et al. Elevated preoperative neutrophil to lymphocyte ratio predicts survival following hepatic resection for colorectal liver metastases. *Eur J Surg Oncol.* 2008; 34: 55-60.