

# Evaluation of the Effects of Thymoquinone on Oxidative Stress in A549 Lung Cancer Cell Line

Reza Farsiabi, MSc, Iraj Khodadadi, PhD, Jamshid Karimi, PhD, Gholamreza Shafiee\*, PhD

Department of Clinical Biochemistry, Medicine School, Hamadan University of Medical Sciences, Hamadan, Iran

Please cite this article as: Farsiabi R, Khodadadi I, Karimi J, Shafiee G. Evaluation of the effects of thymoquinone on oxidative stress in A549 lung cancer cell line. Middle East J Cancer. 2023;14(2):231-40. doi: 10.30476/mejc.2021.92306.1653.

## Abstract

**Background:** Thymoquinone (TQ), an active part of *Nigella sativa*, has been reported as an anticancer agent. This study aimed to evaluate different anticancer effects of TQ on oxidative stress markers and Peroxiredoxin 4 (P4) in lung cancer A549 cell line.

**Method:** In this experimental study, we used TQ concentrations to treat lung A549 cells and determined cell viability by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay at 12, 24, and 48 h times. The IC<sub>50</sub> concentration of TQ was found with MTT assay. We studied the total antioxidant capacity (TAC) and total oxidant status (TOS) using the manual assay, and analyzed catalase (CAT), superoxide dismutases (SOD), and glutathione peroxidase (GPx) activity using the enzyme-linked immunoassay (ELISA) kits. Moreover, the concentration of peroxiredoxin-4 (PRXD4) was measured with the ELISA Kit.

**Results:** The IC<sub>50</sub> of TQ for A549 cells was calculated to be 40 μM concentration for 24 h of incubation. TAC index significantly decreased in the treated cells compared with the controls ( $P = 0.05$ ), whereas TOS and PRXD4 levels showed a significant increase ( $P = 0.05$ ). Additionally, the results showed that the CAT, SOD, and GPx activity enzymes significantly decreased in 20, 40, and 60 μM TQ in comparison with the control cells ( $P = 0.05$ ).

**Conclusion:** TQ has significant inhibitory effects on A549 cells and could be utilized in novel therapy not only for lung cancer, but also for other tumors.

**Keywords:** Thymoquinone, Oxidative stress, Peroxiredoxin 4, Lung neoplasms

## Corresponding Author:

Gholamreza Shafiee, PhD  
Department of Clinical Biochemistry, Medicine School, Nutrition Health Research Center, Hamadan University of Medical Sciences, Hamadan, Iran  
Tel: +98 81 38380462  
Fax: +9881 38380208  
Email: g\_r\_shafiee@yahoo.com

## Introduction


According to the World Health Organization in 2018, lung cancer is the most prevalent type of cancer worldwide and has the highest

mortality rate (1.76 million deaths) compared with other malignant tumors. Lung adenocarcinoma is more common in people over the age of 50, and smoking is the most

effective risk factor for this disease.<sup>1</sup> Lung cancer is classified into two main groups: small cell lung cancer (SCLC) and non-small cell lung cancer (NSCLC); the latter accounts for 85% of cases. A549 Lung cancer cell line is commonly used as an NSCLC cell line in drug discovery and basic research.<sup>2</sup> These cancerous epithelial cells were obtained from a 58-year-old Caucasian male patient.<sup>3</sup> Oxidative stress is a major feature of tissue and cell damage. Several studies have shown the importance of antioxidant and pro-oxidant markers in various cancers.<sup>4</sup> Furthermore, the beneficial role of antioxidant rich diets in the risk of carcinoma diseases has been proven. Oxidative stress occurs when there is an imbalance between the antioxidant /oxidant systems and the cell is unable to neutralize pro-oxidants and free radicals.<sup>1</sup> Reactive oxygen species (ROS) are the main class of radicals produced in cells, which include the hydroxyl, perhydroxyl, peroxy, and superoxide radicals along with non-free radical species, such as singlet oxygen and hydrogen peroxide.<sup>5</sup> Recent studies have shown conflicting results regarding the role of ROS in cancers. On one hand, increase in ROS induces apoptosis<sup>6, 7</sup> by activating caspases and suppressing cancer by inactivating the STAT3 / JAK2 pathway in several types of cancer.<sup>7-10</sup> On

the other hand numerous research findings have indicated that elevated ROS levels may lead to mutagenesis through DNA damage, especially if the accumulation of DNA damage is combined with an incomplete repair system or a deficient apoptotic pathway.<sup>11</sup> The pro-oxidants are constantly produced in living cells; accordingly, certain antioxidant defense systems are needed to counteract them and prevent oxidative stress. These defense systems contain antioxidants, such as vitamin E, vitamin C, and glutathione (GSH) (reduction form) in addition to (antioxidant) enzymes, such as catalase (CAT), glutathione peroxidase (GPx), superoxide dismutase (SOD), and peroxiredoxin 4 (PRXD4).<sup>12, 13</sup> SOD enzyme modulates cellular oxidative stress by scavenging superoxide radicals and converting them to hydrogen peroxide. Afterwards, GPx and CAT eliminate the hydrogen peroxide through oxidation of GSH and producing water, respectively.<sup>14</sup> In addition to CAT and GPx, the Prxs protein family converts H<sub>2</sub>O<sub>2</sub> to water and harmless metabolites; among these PRXDs, PRXD4 plays an important role in tumor metastasis and progression in lung adenocarcinoma.<sup>13</sup>

Thymoquinone (TQ), the bioactive compound of *Nigella sativa* (black seed oil), was first extracted by El-Dakhkhany and is mostly



**Figure 1.** The effect of thymoquinone on A549 cell viability. Exposure of A549 cells to different concentrations of thymoquinone for 24 and 48 h decreased cell viability in a dose-dependent manner. The incubation of the cells with thymoquinone for 24 h exhibited a greater decline in the cell viability compared with longer incubation times.

known as an antioxidant, antineoplastic, and antiinflammatory agent in both *in vitro* and *in vivo* xenograft models.<sup>15</sup> The antitumor effects of TQ on decreasing cell viability, inducing apoptosis, inhibiting cell proliferation, and generation of ROS have been proven by numerous papers, yet its central mechanism of action has not been fully understood.<sup>16-18</sup> In addition, there are contradictory reports about the dual effect of TQ on oxidative stress in different cancers. This led us to design a (novel) study to investigate a group of oxidative stress markers in order to attain insights into the effect of TQ on oxidative stress on lung cancer cells and possibly elucidate the antitumor mechanism of TQ. We assessed oxidative stress by measuring the total oxidative status (TOS), total antioxidant capacity (TAC), PRXD4 levels, CAT, GPx, and SOD enzymes activities.

## Material and Methods


### Cell culture

This work was designed as an experimental study. We purchased human A549 lung epithelial (human non-small lung cancer cell line) cancer cells from Pasteur Institute cell culture collection (Tehran, Iran). A549 cells have a high proliferation rate and long lifespan and maintain their phenotype in culture. RPMI1640 culture media, FBS, 0.25%

Trypsin-EDTA solution, and phosphate-buffered saline (PBS) tablets were obtained from Life Technologies (Life Technologies Ltd., Paisley-UK) and Sigma-Aldrich Company (Sigma-Aldrich Co., Steinheim, Germany) supplied DMSO, penicillin, streptomycin, and TQ. A549 cells were cultured in 25-cm<sup>2</sup> flasks in RPMI1640 medium containing 10% fetal bovine serum and 1% penicillin-streptomycin (100 units/ml penicillin and 100 µg/ml streptomycin). The cultured cells were grown at 37°C in 5% CO<sub>2</sub> humidified air.

### MTT assay

The cell viability was evaluated with MTT (3-[4, 5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) assay. Briefly, the cells were seeded at a 1×10<sup>4</sup>/well density in 96 well plates and incubated at 37°C with 5% CO<sub>2</sub>. We carried out the treatment with different concentrations of TQ (0, 20, 40, 60, 80, 100 µM) and incubated the cells for 24 and 48h. Subsequently, 10µl MTT (5mg/ml) was added to each well and the plate was incubated for 4h at 37°C in the dark so that formazan could be formed by mitochondrial succinate dehydrogenase. The content of wells was replaced with 100µL DMSO and the absorbance was measured at 570 nm employing ELISA plate reader (Sunrise Tecan, Switzerland). The inhibitory rates were calculated via the



**Figure 2.** The effects of thymoquinone on the Prx4 concentration in A549 cells. The cells were treated with 0, 20, 40, and 60 mM of thymoquinone; the lowest level of Prx4 concentration was observed in the cells treated with 40 µM thymoquinone. \* is a significant sign with the control. # is a significant sign with the 20 µM thymoquinone. Mean ± SD is considered in all the diagrams.  $P < 0.05$  was considered as the level of significance.

following formula:

$$\text{Inhibitory rates (IR)} = (1 - (\text{OD}_{\text{treatment}} / \text{OD}_{\text{control}})) \times 100$$

The  $\text{IC}_{50}$  was calculated using the following formula according to the inhibitory rate:

$$\text{IC}_{50} = (50\% - \text{Low}_{\text{inh}}\%) / (\text{High}_{\text{inh}}\% - \text{Low}_{\text{inh}}\%) \times (\text{High}_{\text{conc}} - \text{Low}_{\text{conc}}) + \text{Low}_{\text{conc}}$$

$\text{Low}_{\text{conc}}$  and  $\text{High}_{\text{conc}}$  were the lowest and highest concentrations of TQ, respectively, and  $\text{Low}_{\text{IR}}$  and  $\text{High}_{\text{IR}}$  were respectively defined as the lowest and highest inhibitory rates.


#### Lysate preparation

The cells were trypsinized and after

centrifugation, the cells were washed with cold PBS twice. A PBS-containing protease inhibitor cocktail was used to prepare cell suspension. The cell lysates were then obtained by repeated freeze-thaw in triplicate. The suspension was centrifuged at 12000 rpm for 15 minutes and the supernatant was collected in the tubes that were stored at  $-20^{\circ}\text{C}$  until further analysis.

#### PRXD4

PRXD4 was measured with the ZellBio-GmbH ELISA Kit (Cat. ZB-12927C-H9648, Ulm Germany) according to the manufacturer's protocol. It was based on colorimetric method



**Figure 3.** The effects of thymoquinone on the TAC and TOS concentrations in A549 cells. The cells were treated with 0, 20, 40, and 60 mM of thymoquinone; in TAC (A) and TOS (B) levels, there was a significant difference in the cells treated with 20, 40, and 60  $\mu\text{M}$  thymoquinone compared to the controls. \* is a significant sign with the control group. # is a significant sign with the 60  $\mu\text{M}$  thymoquinone. Mean  $\pm$  SD is considered in all the diagrams.  $P < 0.05$  was considered as the level of significance.

TAC: Total antioxidant capacity; TOS: Total oxidant status

and the basis of the measurement was a non-competitive sandwich assay. The standard range for Prx4 concentration was 1.25-20 ng/ml.

#### TAC

TAC measurement was performed utilizing FRAP manual method. It assessed the ability of antioxidant compounds of sample to reduce  $\text{Fe}^{3+}$ -TPTZ complex to  $\text{Fe}^{2+}$ -TPTZ. The reaction mixture comprised 300  $\mu\text{L}$  FARP reagent (acetate buffer 300 mM, pH 3.6, TPTZ 10 mM and ferric chloride 20 mM) and 10  $\mu\text{L}$  of the samples was pipetted into 96 well plates and incubated for 10 min in the dark at 37°C. The concentrations of standards (1M  $\text{FeSO}_4$ ) and the samples were measured with Spectrophotometer (Bell-Italy) at a wavelength of 593 nm. The concentration of each sample was then determined according to the standard curve.<sup>19</sup>

#### TOS

TOS was evaluated with ferric-xylenol orange (FOX) assay based on the oxidation of ferrous ions to ferric ions, which results in a colored ferric-xylenol orange complex. The reaction mixture included 10  $\mu\text{L}$  of the sample and 190  $\mu\text{L}$  of the FOX reagent (250  $\mu\text{M}$  ferrous ion, 150  $\mu\text{M}$  xylenol orange, 100 mM sorbitol and 25 mM  $\text{H}_2\text{SO}_4$ , pH was 1.8), which was inserted into 96 well plates.  $\text{H}_2\text{O}_2$  and PBS were applied as standards and blank, respectively. The absorbance at 560 nm was read after 10 min of incubation in the room temperature.<sup>20</sup> Finally, the oxidative stress index (OSI) was defined as the percentage ratio of TOS to TAC.

#### GPx enzyme activity

GPx activity was determined with a ready-to-use kiazist kit (Tehran, Iran) based on the reduction in  $\text{H}_2\text{O}_2$  by glutathione oxidation, which acts as an electron donor. GSH enzyme was converted to GSSG, and the remaining GSH in the reaction medium could regenerate DTNB and produce a yellow color absorbed at 412 nanometers. Color production was inversely related to enzyme activity. Finally, the GPx activity was reported as U/mg protein.

#### SOD enzyme activity

SOD activity was assayed with a ready-to-use kiazist kit (Tehran, Iran). In this method,

superoxide anion was utilized to convert resazurin to resorufin and to produce chromogen. Ultimately, the chromogen was measured in terms of calorific value at 420 nm and the results were reported as U/mg protein.

#### CAT enzyme activity

Catalase activity was measured using the previously described spectrophotometric method by Hadwan and Abed.<sup>21</sup> Briefly, 100  $\mu\text{L}$  of the samples was added and incubated with 1 ml reagent (20 mM  $\text{H}_2\text{O}_2$  in 50 mM sodium phosphate buffer, pH 7.4) at 37°C for 3 min. Afterwards, 4  $\mu\text{L}$  of ammonium molybdate (32.4 mM) was added to stop the reaction. The absorbance of the obtained yellowish mixture could be measured at 374 nm. Catalase activity was calculated via the following formula:

$$\text{Catalase Activity kU} = 2.303/t * [\log S^0/S-M] * V_t/V_s$$

#### Data analysis

Data was analyzed employing the statistical software SPSS 16 (SPSS Inc. Chicago, USA) and t-test was performed to compare the differences between the groups. One-way ANOVA, followed by Tukey-Kramer multiple comparisons test, was utilized to analyze the differences between the groups. The values are given as mean  $\pm$  SD. The significance level of all the statistical tests was considered to be at 0.05.

## Results

### TQ inhibited viability and changed the morphology of A549 lung cancer cells

As shown in figure 1, A549 lung cancer cell viability decreased significantly after 24 h of incubation with TQ in a dose-dependent manner ( $P < 0.05$ ).  $\text{IC}_{50}$  of TQ for A549 cells was calculated as 40  $\mu\text{M}$  concentration; 20 and 60  $\mu\text{M}$  concentrations were also used in further experiments as low and high doses, respectively. TQ treatment changed the morphology of the cells. The morphology of A549 adherent epithelial cells changed. They became round in shape and were suspended in their media dose-dependently.

#### PRXD4


Figure 2 represents a significant dose-dependent decrease in A549 lung cancer cells in

PRXD4 concentration after exposure of the cells to 20, 40, and 60 mM. The most PRXD4 decrease was observed in 40  $\mu\text{M}$  TQ dose ( $P < 0.05$ ). Moreover, the cells treated by 40 and 60  $\mu\text{M}$  TQ showed a significant difference with those treated

via a 20- $\mu\text{M}$  dose ( $P < 0.05$ ).

#### TAC and TOS

The findings showed a significant change in TAC index in the cells treated with 20, 40, and 60 mM TQ compared with that of the controls



**Figure 4.** The effects of thymoquinone on the CAT, SOD, and GPx activities in A549 cells. The cells were treated with 0, 20, 40, and 60 mM of thymoquinone; in CAT (A), SOD (B), and GPx (C) activities, there was a significant difference between the cells treated with 20, 40, and 60  $\mu\text{M}$  thymoquinone compared with the controls. \* is a significant sign with the control. # is a significant sign with the 60  $\mu\text{M}$  thymoquinone. \* is a significant sign with the 20 and 40  $\mu\text{M}$  thymoquinone. Mean  $\pm$  SD is considered in all the diagrams.  $P < 0.05$  was considered as the level of significance.

CAT: Catalase; SOD: Superoxide dismutase; GPx: Glutathione peroxidase

( $P = 0.05$ ), as shown in figure 3A. Additionally, the cells treated with 20 and 40  $\mu\text{M}$  TQ had a significant difference with those treated with a 60- $\mu\text{M}$  dose ( $P < 0.05$ ). The result concerning TOS levels indicated a difference between 20, 40, and 60  $\mu\text{M}$  concentrations of TQ with the controls ( $P = 0.05$ ), as demonstrated in figure 3B. The cells treated with 20 and 40  $\mu\text{M}$  TQ had a significant difference with those treated with a 60- $\mu\text{M}$  dose ( $P < 0.05$ ).

#### *CAT, SOD, and GPx*

The results of the study of CAT enzyme activity implied that the activity of this enzyme significantly decreased dose-dependently in 20, 40, and 60  $\mu\text{M}$  TQ compared with that of the controls ( $P = 0.05$ ) (Figure 4A). The results regarding SOD enzyme activity revealed that the activity decreased dose-dependently in 20, 40, and 60  $\mu\text{M}$  TQ compared with that of the controls ( $P = 0.05$ ) (Figure 4B). As shown in figure 4C, there was a significant reduction in the amount of GPx enzyme activity in the treated cells compared with the controls. Moreover, there was a significant difference in the cells treated with 20 and 40  $\mu\text{M}$  compared with those treated with 60  $\mu\text{M}$  TQ concentration ( $P < 0.05$ ).

### Discussion

Even though there are a number of reports on oxidant/antioxidant properties of TQ on lung cancer, this is the first study to investigate the potential effects of TQ on oxidative stress parameters in A549 adenocarcinoma lung cancer cell line. Our findings demonstrated that TQ exposure induced oxidative stress by increasing TOS and decreasing PRDX4, the activity of GPx, SOD, CAT, and TAC in A549 cells. We revealed that the most pro-oxidant effects of TQ were seen in low concentrations. The imbalance between oxidants and antioxidant systems leads to oxidative stress. TOS and TAC are two reliable and precise indicators, which are mostly used to determine oxidative status.<sup>22, 23</sup> Our results indicated that TQ induced oxidative stress by elevating TOS and declining TAC significantly in A549 cells. Yih-Fang Liou et al. suggested the antiproliferative and anticancer properties of TQ due to reactive oxygen

species/superoxide generation. Their results showed the production of ROS in 786-O-SI3 renal cancer cells and BFTC-909 transitional cancer cells.<sup>24</sup> In addition, Ji Eun Park et al. conducted a study to provide a mechanism for inducing apoptosis by TQ in A431 epidermoid carcinoma cells. They indicated that exposure to the TQ prompts cells into the oxidative status and induces the generation of ROS.<sup>25</sup> Furthermore, the role of TQ in ROS induction in MDA-MB-231 and MCF-7 breast cancer cell lines and xenograft mouse model has been reported.<sup>25</sup> EM Dergarabetian et al. showed that TQ induced apoptosis in malignant T-cells through ROS generation.<sup>26</sup> Despite the presented data and mentioned studies, there are reports considering TQ as an antioxidant agent. The antioxidant property of TQ is supported by the decrease in TOS and increase in TAC in Ehrlich acid mouse solid tumor models and mouse mesenchymal stem cells, respectively.<sup>26, 27</sup> SOD, GPx, and CAT are the main antioxidant enzymes that play a crucial role in reducing oxygen radicals and cell oxidative status. SOD converts superoxide and single-chain radicals to an oxygen molecule and hydrogen peroxide. Finally, GPx and CAT remove the produced  $\text{H}_2\text{O}_2$  to avoid converting it to harmful substances, such as hydroxyl radicals by Fenton reaction.<sup>22</sup> Our findings showed that SOD, GPx, and CAT activities significantly reduced in the treated cells compared with those in the untreated ones. Certain investigations have reported that TQ results in an increase in superoxide anion.<sup>24, 28</sup> Increased superoxide anion in those studies is a result of the reduction in SOD activity, which leads to a decrease in  $\text{H}_2\text{O}_2$  production. Therefore, the lack of enough  $\text{H}_2\text{O}_2$  as the substrate of GPx and CAT possibly decreases their activity. It has been also suggested that reduction in SOD activity and  $\text{H}_2\text{O}_2$  for the reason behind increased superoxide anion. It is predicted that CAT and GPx activity decrease in response to  $\text{H}_2\text{O}_2$  reduction. In contrast with our observations, it has been demonstrated that TQ can attenuate oxidative stress by increasing SOD, GPx, and CAT after partial hepatectomy on Ischemic reperfusion liver of rats. TQ antioxidant

effects were explained by its ability to upregulate antioxidant genes and scavenge free radicals.<sup>29</sup>

PRXD4 is a member of peroxiredoxin family proteins, which converts alkyl hydroperoxides and H<sub>2</sub>O<sub>2</sub> to alcohol and water through the use of thiol-containing donor molecules, such as GSH and thioredoxine (Trx).<sup>30</sup> Our observations revealed a decrease in PRDX4 after 24 h of TQ exposure. It has been reported that TQ consumption results in GSH depletion in mice tissues, like the liver, kidneys, and heart.<sup>31</sup> In addition, it has been shown that GSH declines in HEp-2 and rat C6 glioma cells following TQ treatment.<sup>28, 32</sup> According to these findings, it could be predicted that decreased PRDX4 might be associated with GSH level.

The obtained data herein revealed that the pro-oxidant effects of TQ declined in a dose-dependent manner. Recently, published papers have provided some evidence indicating that TQ induces oxidative stress and ROS generation, specifically at low doses (20-50 µM).<sup>28</sup> They have also reported that 40 µM TQ can increase ROS in various cell lines.<sup>33</sup> The ROS production has been attenuated in the high concentration of TQ.<sup>34</sup> According to these pieces of evidence, it could be suggested that the oxidant/antioxidant effects of TQ depend on its concentration. It seems as though the conflicting effect of TQ on oxidative status possibly depends on its concentration. Contradictory reports concerning pro-oxidant/antioxidant ability of TQ are attributed to its reduction or oxidation (reduction-oxidation) form. Quinones have been known as a redox-cycling compound that can undergo one- or two-electron reductions by cellular reductases. One-electron reduction yields semiquinone that can act as a pro-oxidant and react with molecular oxygen (O<sub>2</sub>) to form hydrogen peroxide and superoxide.<sup>28, 33</sup> This reaction catalyzes NADH-cytochrome b5 reductase, NADPH-cytochrome P450 reductase, NADH: ubiquinone oxidoreductase. Two-electron reduction yields hydroquinone which is catalyzed by NADH-Quinone oxidoreductase 1 (NQO1) and acts as an antioxidant.<sup>28, 34</sup> According to the induced oxidative stress by TQ in our study, it seems as

if in A549 cell lines, the predominant form of quinones is semiquinone. However, further research is recommended to confirm this result. Investigating cellular reductases, such as NQO1 and NADPH-cytochrome P450 reductase involved in TQ reduction, may be conducive to finding the mechanism of TQ in inducing oxidative stress in A549 cells. In addition, using PCR to investigate the related genes (GPx, CAT, SOD) and signaling pathways (STAT3 / JAK2) could provide valuable information about the effects of TQ on oxidative stress.

## Conclusion

In conclusion, our study shed light on the fact that TQ induced oxidative stress in A549 lung cancer cells. TQ treatment increased TOS and decreased TAC, PRDX4, SOD, GPx, and CAT activity and the most pro-oxidant effects belonged to low concentrations. These results suggested that the mechanism of TQ as a potential anticancer agent may be owing to its pro-oxidant ability in A549 lung cancer cells. Furthermore, using the PCR array method was shown to provide valuable information about the effects of TQ on oxidative stress and other signaling pathways involved in lung cancer.

## Acknowledgments

The authors wish to acknowledge the financial support of Hamadan University of Medical Sciences (Project NO: 990216748).

## Conflict of Interest

None declared.

## References

1. Aybastýer Ö, Dawbaa S, Demir C, Akgün O, Ulukaya E, Arý F. Quantification of DNA damage products by gas chromatography tandem mass spectrometry in lung cell lines and prevention effect of thyme antioxidants on oxidative induced DNA damage. *Mutat Res.* 2018;808:1-9. doi: 10.1016/j.mrfmmm.2018.01.004.
2. Chan DC, Earle KA, Zhao TL, Helfrich B, Zeng C, Baron A, et al. Exisulind in combination with docetaxel inhibits growth and metastasis of human lung cancer and prolongs survival in athymic nude rats with orthotopic lung tumors. *Clin Cancer Res.*

- 2002;8(3):904-12.
3. Samarghandian S, Azimi-Nezhad M, Farkhondeh T. Thymoquinone-induced antitumor and apoptosis in human lung adenocarcinoma cells. *J Cell Physiol.* 2019;234(7):10421-31. doi: 10.1002/jcp. 27710.
  4. Jrah-Harzallah H, Ben-Hadj-Khalifa S, Almawi WY, Maaloul A, Houas Z, Mahjoub T. Effect of thymoquinone on 1,2-dimethyl-hydrazine-induced oxidative stress during initiation and promotion of colon carcinogenesis. *Eur J Cancer.* 2013;49(5): 1127-35. doi: 10.1016/j.ejca.2012.10.007.
  5. Quiñonez-Flores CM, González-Chávez SA, Del Río Nájera D, Pacheco-Tena C. Oxidative stress relevance in the pathogenesis of the rheumatoid arthritis: A systematic review. *Biomed Res Int.* 2016;2016: 6097417. doi: 10.1155/2016/6097417.
  6. Cockfield JA, Schafer ZT. Antioxidant defenses: A context-specific vulnerability of cancer cells. *Cancers.* 2019;11(8):1208. doi.org/10.3390/cancers11081208.
  7. Sun Q, Lu NN, Feng L. Apigenin inhibits gastric cancer progression through inducing apoptosis and regulating ROS-modulated STAT3/JAK2 pathway. *Biochem Biophys Res Commun.* 2018;498(1):164-70. doi: 10.1016/j.bbrc.2018.02.009.
  8. He H, Zhuo R, Dai J, Wang X, Huang X, Wang H, et al. Chelerythrine induces apoptosis via ROS-mediated endoplasmic reticulum stress and STAT3 pathways in human renal cell carcinoma. *J Cell Mol Med.* 2020;24(1):50-60. doi: 10.1111/jcmm.14295.
  9. Chae IG, Kim DH, Kundu J, Jeong CH, Kundu JK, Chun KS. Generation of ROS by CAY10598 leads to inactivation of STAT3 signaling and induction of apoptosis in human colon cancer HCT116 cells. *Free Radic Res.* 2014;48(11):1311-21. doi: 10.3109/10715762.2014.951838.
  10. Miao Z, Yu F, Ren Y, Yang J. d,l-Sulforaphane induces ROS-dependent apoptosis in human glioblastoma cells by inactivating STAT3 signaling pathway. *Int J Mol Sci.* 2017;18(1):72. doi: 10.3390/ijms18010072.
  11. Prasad S, Gupta SC, Tyagi AK. Reactive oxygen species (ROS) and cancer: Role of antioxidative nutraceuticals. *Cancer Lett.* 2017;387:95-105. doi: 10.1016/j.canlet.2016.03.042.
  12. Backos DS, Franklin CC, Reigan P. The role of glutathione in brain tumor drug resistance. *Biochem Pharmacol.* 2012;83(8):1005-12. doi: 10.1016/j.bcp.2011.11.016.
  13. Mishra M, Jiang H, Wu L, Chawsheen HA, Wei Q. The sulfiredoxin-peroxiredoxin (Srx-Prx) axis in cell signal transduction and cancer development. *Cancer Lett.* 2015;366(2):150-9. doi: 10.1016/j.canlet. 2015.07. 002.
  14. Arab Sadeghabadi Z, Abbasalipourkabir R, Mohseni R, Ziamajidi N. Investigation of oxidative stress markers and antioxidant enzymes activity in newly diagnosed type 2 diabetes patients and healthy subjects, association with IL-6 level. *J Diabetes Metab Disord.* 2019;18(2):437-43. doi: 10.1007/s40200-019-00437-8.
  15. Gali-Muhtasib H, Roessner A, Schneider-Stock R. Thymoquinone: a promising anti-cancer drug from natural sources. *Int J Biochem Cell Biol.* 2006;38(8):1249-53. doi: 10.1016/j.biocel.2005.10.009.
  16. Ulasli SS, Celik S, Gunay E, Ozdemir M, Hazman O, Ozyurek A, et al. Anticancer effects of thymoquinone, caffeic acid phenethyl ester and resveratrol on A549 non-small cell lung cancer cells exposed to benzo(a)pyrene. *Asian Pac J Cancer Prev.* 2013;14(10):6159-64. doi: 10.7314/apjcp.2013. 14.10.6159.
  17. Shafiee G, Saidijam M, Tayebinia H, Khodadadi I. Beneficial effects of genistein in suppression of proliferation, inhibition of metastasis, and induction of apoptosis in PC3 prostate cancer cells. *Arch Physiol Biochem.* 2020:1-9. doi: 10.1080/13813455.2020. 1717541.
  18. Shafiee G, Saidijam M, Tavilani H, Ghasemkhani N, Khodadadi I. Genistein induces apoptosis and inhibits proliferation of HT29 colon cancer cells. *Int J Mol Cell Med.* 2016;5(3):178-91.
  19. Benzie IF, Szeto YT. Total antioxidant capacity of teas by the ferric reducing/antioxidant power assay. *J Agric Food Chem.* 1999;47(2):633-6. doi: 10.1021/jf9807768.
  20. Erel O. A new automated colorimetric method for measuring total oxidant status. *Clin Biochem.* 2005;38(12):1103-11. doi: 10.1016/j.clinbiochem. 2005.08.008.
  21. Hadwan MH, Abed HN. Data supporting the spectrophotometric method for the estimation of catalase activity. *Data Brief.* 2015;6:194-9. doi: 10.1016/j.dib.2015.12.012.
  22. Bahmani M, Ziamajidi N, Hashemnia M, Abbasalipourkabir R, Salehzadeh A. Human umbilical cord blood mesenchymal stem cell conditioned medium (hMSC-CM) improves antioxidant status in carbon tetrachloride-induced oxidative damage in rat. *Iran J Sci Technol Trans Sci.* 2020;44(5):1327-35. doi:10.1007/s40995-020-00944-x.
  23. Şen V, Bozkurt M, Söker S, Ece A, Güneş A, Uluca Ü, et al. The effects of pomegranate and carvacrol on methotrexate-induced bone marrow toxicity in rats. *Clin Invest Med.* 2014;37(2):E93-E101. doi: 10.25011/cim.v37i2.21091.
  24. Liou YF, Chen PN, Chu SC, Kao SH, Chang YZ, Hsieh YS, et al. Thymoquinone suppresses the proliferation of renal cell carcinoma cells via reactive oxygen species-induced apoptosis and reduces cell stemness. *Environ Toxicol.* 2019;34(11):1208-20. doi: 10.1002/tox.22822.
  25. Park JE, Kim DH, Ha E, Choi SM, Choi JS, Chun KS, et al. Thymoquinone induces apoptosis of human

- epidermoid carcinoma A431 cells through ROS-mediated suppression of STAT3. *Chem Biol Interact.* 2019;312:108799. doi: 10.1016/j.cbi.2019.108799.
26. Meral I, Pala M, Akbas F, Ustunova S, Yildiz C, Demirel MH. Effects of thymoquinone on liver miRNAs and oxidative stress in Ehrlich acid mouse solid tumor model. *Biotech Histochem.* 2018;93(4): 301-8. doi: 10.1080/10520295.2018.1437472.
  27. Rezaei N, Sardarzadeh T, Sisakhtnezhad S. Thymoquinone promotes mouse mesenchymal stem cells migration in vitro and induces their immunogenicity in vivo. *Toxicol Appl Pharmacol.* 2020;387:114851. doi: 10.1016/j.taap.2019.114851.
  28. Krylova NG, Drobysh MS, Semenkova GN, Kulahava TA, Pinchuk SV, Shadyro OI. Cytotoxic and antiproliferative effects of thymoquinone on rat C6 glioma cells depend on oxidative stress. *Mol Cell Biochem.* 2019;462(1-2):195-206. doi: 10.1007/s11010-019-03622-8.
  29. Bouhlel A, Bejaoui M, Ben Mosbah I, Hadj Abdallah N, Ribault C, Viel R, et al. Thymoquinone protects rat liver after partial hepatectomy under ischaemia/reperfusion through oxidative stress and endoplasmic reticulum stress prevention. *Clin Exp Pharmacol Physiol.* 2018. doi: 10.1111/1440-1681.12961.
  30. Fujii J, Ikeda Y. Advances in our understanding of peroxiredoxin, a multifunctional, mammalian redox protein. *Redox Rep.* 2002;7(3):123-30. doi: 10.1179/135100002125000352.
  31. Badary OA, AlShabanah OA, Nagi MN, Al-Bekairi AM, Elmazar M. Acute and subchronic toxicity of thymoquinone in mice. *Drug Development Research.* 1998;44(2-3):56-61. doi.org/10.1002/(SICI)1098-2299(199806/07)44:2/3<56::AID-DDR2>3.0.CO;2-9.
  32. Rooney S, Ryan MF. Modes of action of alpha-hederin and thymoquinone, active constituents of *Nigella sativa*, against HEP-2 cancer cells. *Anticancer Res.* 2005;25(6B):4255-9.
  33. El-Najjar N, Chatila M, Moukadem H, Vuorela H, Ocker M, Gandesiri M, et al. Reactive oxygen species mediate thymoquinone-induced apoptosis and activate ERK and JNK signaling. *Apoptosis.* 2010;15(2):183-95. doi: 10.1007/s10495-009-0421-z.
  34. Ashour AE, Abd-Allah AR, Korashy HM, Attia SM, Alzahrani AZ, Saquib Q, et al. Thymoquinone suppression of the human hepatocellular carcinoma cell growth involves inhibition of IL-8 expression, elevated levels of TRAIL receptors, oxidative stress and apoptosis. *Mol Cell Biochem.* 2014;389(1-2):85-98. doi: 10.1007/s11010-013-1930-1.