

The Paradigm of Technological Innovations from Open to Scarless Surgery for Colorectal Cancer

Salman Yousuf Guraya

Clinical Sciences Department, College of Medicine, University of Sharjah, UAE

The global burden of colorectal cancer (CRC) and therapeutic evolutions

The estimated incidence of colorectal cancer (CRC) is expected to increase by 60% with over 2.2 million new cases and 1.1 million deaths by 2030.^{1,2} In keeping with this global escalation of CRC, the pedagogy of curative therapies has also been revolutionized. Innovations in bioengineering technologies show tremendous progress in the surgical treatments of CRC with a significant impact on postoperative recovery time, the amount of surgical trauma, and cancer-related morbidity and survival.³ The 4-year disease-free survival for patients with stages I-III CRC has improved to 85.5% following total mesorectal excision.⁴ The following sections of this paper shed light on the surgical developments for CRC that have accounted for less complications and morbidity, rapid recovery, and improved survival.

Surgical management of colorectal cancer (CRC) by laparoscopic and robotic assisted laparoscopic surgery

The management of CRC primarily involves a multi-disciplinary approach that derives services from a number of disciplines - surgery, oncology, radiology, nutrition, and social care.⁵ The main reason for integrating these disciplines is driven by the fact that the disease quite often presents at an advanced stage which necessitates neo(adjuvant) therapy and nutritional support during the perioperative period.⁶ Traditionally, the mainstay of CRC treatment is surgical therapy that carries the promise of cure even in the presence of metastases. Before the advent of laparoscopy, all surgical therapies were performed by an open technique with a high incidence of anastomotic leakage,⁷ wound infection, adhesions and incisional hernia,⁸ longer hospital stay, and high morbidity.⁹ In the 1990s, laparoscopic surgery revolutionized the landscape of surgical therapies in all disciplines,

♦Corresponding Author:

Salman Yousuf Guraya, FRCS,
Masters MedEd
Professor of Surgery, Vice
Dean, College of Medicine,
Clinical Sciences Department,
University of Sharjah, UAE
Email: salmanguraya@gmail.com

including CRC. However, the learning curve for CRC has remained long and tedious.¹⁰ Nevertheless, the cutting-edge developments in surgical techniques as well as the state-of-the-art programs overcame this shortcoming to some extent and offer well-structured accredited platforms for surgical training.¹¹ The educational reforms that frame the entire training of medical graduates in the work-place environment can potentially enrich the learning avenues for those students who intend to pursue a surgical career.¹² The outright benefits of laparoscopic surgery for CRC are far ranging and include shorter hospital stay, less blood loss, high lymph node yield, less postoperative pain, and rapid return of bowel function compared to the open technique.^{13,14} A study by Schwenk et al. has shown a lower incidence of postoperative complications (18.2%) in patients who underwent laparoscopic resection for CRC compared with open surgery (23%).¹⁵

The two-dimensional video imaging, unstable assistant-held camera platform, limited range of motion, and poor surgical ergonomics undermine the effectiveness and feasibility of laparoscopic surgery for CRC.¹⁶ Although there is convincing evidence of enhanced convalescence and low morbidity by laparoscopic surgery compared to the open technique,^{3,17} some reports have shown tumor recurrence at the port sites, which has challenged the oncological safety of laparoscopic surgery.^{18,19} Unfortunately, the large incision to retrieve the surgical specimen through the abdominal wall defeats the main purpose of using small incisions by laparoscopic surgery. The three-dimensional video imaging, enormous range of motion, and a very stable camera platform by robotic surgery has overcome some of the pitfalls witnessed during laparoscopic surgery.²⁰ Interestingly, research has not shown a significant difference in the 5-year disease-free survival as well as for locoregional recurrence between laparoscopic and robotic-assisted laparoscopic surgery.²¹ Also, the challenges of wound seeding by cancer cells and large incision for specimen retrieval have been reported with the use of robotic assisted laparoscopic surgery for CRC.^{22,23}

Natural orifice transluminal endoscopic surgery (NOTES)

The innovations in bioengineering have recently introduced natural orifice transluminal endoscopic surgery (NOTES) that allows surgical access within the abdominal or thoracic cavities through a natural orifice – either the mouth, anus, urethra, or vagina.²⁴ Although the overarching benefit of NOTES remains the scarless surgery, this surgical strategy requires visceral perforation and closure such as transcolonic, transvaginal, and transgastric approaches. With NOTES, the concept of scarless surgery has been materialized, recovery is quick even after major surgical procedures, and there is significantly less post-procedural pain.²⁵ ‘Hybrid’ NOTES entails the surgical entities that requires some type of transabdominal assistance. In some situations, the majority of the procedure is conducted using transabdominal instruments, whereas other surgeries are predominantly transluminal but need transabdominal assistance.²⁶ The transvaginal approach has been successfully used for large bowel cancers by intracorporeal anastomosis and transvaginal specimen extraction,²⁷ nephrectomy,²⁸ partial gastrectomy,²⁹ and cholecystectomy.³⁰ Saad and Hosogi have adopted a new transanal approach that included a standard laparoscopic left colon resection and NOTES by introducing a resectoscope into the rectum, where they removed the specimen through the same route.³¹ Lastly, Targarona et al. adopted a transgastric approach for splenic disorders where spleen was extracted transvaginally.³²

Challenges of scarless surgery by natural orifice transluminal endoscopic surgery (NOTES) and the future

Natural orifice transluminal endoscopic surgery marks a landmark leap in medicine by introducing scarless surgery that uses natural orifices. In addition, benefits of NOTES include better cosmesis, less post-operative pain with early recovery and a short hospital stay. However, despite its great promise, the future of NOTES remains unclear due to profound ethical issues,

procedural challenges and difficulties, and an extremely long learning curve. The pressing technical challenge in NOTES is the narrow operative field with limited maneuverability allowed by the natural orifice. A much higher cost and a long learning curve for NOTES compared to laparoscopic assisted surgery for CRC have also been reported.³³ The cost effectiveness of NOTES is the biggest hurdle against its global application. Compromised sterilization and insecure closure of the penetrations of the stomach and colon pose the greatest challenge to the use of transgastric or transcolonic NOTES.³⁴ The safety and efficacy of this surgical development needs to be established by further animal and human studies in clinical trials. Once proven safe and feasible, NOTES carries great promise for scarless surgery in the modern era.

References

1. Arnold M, Sierra MS, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global patterns and trends in colorectal cancer incidence and mortality. *Gut*. 2017;66(4):683-91.
2. Singh KE, Taylor TH, Pan CG, Stamos MJ, Zell JA. Colorectal cancer incidence among young adults in California. *J Adolesc Young Adult Oncol*. 2014;3(4):176-84.
3. Lacy AM, García-Valdecasas JC, Delgado S, Castells A, Taurá P, Piqué JM, et al. Laparoscopy-assisted colectomy versus open colectomy for treatment of non-metastatic colon cancer: a randomised trial. *Lancet*. 2002;359(9325):2224-9.
4. Bertelsen CA, Neuenschwander AU, Jansen JE, Wilhelmsen M, Kirkegaard-Klitbo A, Tenma JR, et al. Disease-free survival after complete mesocolic excision compared with conventional colon cancer surgery: a retrospective, population-based study. *Lancet Oncol*. 2015;16(2):161-8.
5. Al-Qahtani MF, Guraya SY. Measuring the attitudes of healthcare faculty members towards interprofessional education in KSA. *Journal of Taibah University Medical Sciences*. 2016;11(6):586-93.
6. Guraya SY. Modern oncosurgical treatment strategies for synchronous liver metastases from colorectal cancer. *Journal of Microscopy and Ultrastructure (JMAU)*. 2013;1(1):1-7.
7. Veldkamp R, Kuhry E, Hop WC, Jeekel J, Kazemier G, Bonjer HJ, et al. Laparoscopic surgery versus open surgery for colon cancer: short-term outcomes of a randomised trial. *Lancet Oncol*. 2005;6(7):477-84.
8. Taylor G, Jayne D, Brown S, Thorpe H, Brown J, Dewberry S, et al. Adhesions and incisional hernias following laparoscopic versus open surgery for colorectal cancer in the CLASICC trial. *Br J Surg*. 2010;97(1):70-8.
9. Colon Cancer Laparoscopic or Open Resection Study Group, Buunen M, Veldkamp R, Hop WC, Kuhry E, Jeekel J, et al. Survival after laparoscopic surgery versus open surgery for colon cancer: long-term outcome of a randomised clinical trial. *Lancet Oncol*. 2009;10(1):44-52.
10. Forgione A, Kislov V, Guraya SY, Kasakevich E, Pugliese R. Safe introduction of laparoscopic colorectal surgery even in remote areas of the world: the value of a comprehensive telementoring training program. *J Laparoendosc Adv Surg Tech A*. 2015;25(1):37-42.
11. Guraya SY, Forgione A, Sampogna G, Pugliese R. The mapping of preferred resources for surgical education: Perceptions of surgical trainees at the Advanced International Minimally Invasive Surgery Academy (AIMS), Milan, Italy. *Journal of Taibah University Medical Sciences*. 2015;10(4):396-404.
12. Guraya SY. Workplace-based assessment; applications and educational impact. *Malays J Med Sci*. 2015;22(6):5-10.
13. Franklin ME, Rosenthal D, Abrego-Medina D, Dorman JP, Glass JL, Norem R, et al. Prospective comparison of open vs. laparoscopic colon surgery for carcinoma. *Dis Colon Rectum*. 1996;39(10 Suppl):S35-46.
14. Fleshman J, Sargent DJ, Green E, Anvari M, Stryker SJ, Beart Jr RW, et al. Laparoscopic colectomy for cancer is not inferior to open surgery based on 5-year data from the COST Study Group trial. *Ann Surg*. 2007; 246(4):655-62; discussion 662.
15. Schwenk W, Haase O, Neudecker J, Müller JM. Short term benefits for laparoscopic colorectal resection. *Cochrane Database Syst Rev*. 2005;(3):CD003145.
16. Ballantyne GH. The pitfalls of laparoscopic surgery: challenges for robotics and telerobotic surgery. *Surg Laparosc Endosc Percutan Tech*. 2002;12(1):1-5.
17. Clinical Outcomes of Surgical Therapy Study Group, Nelson H, Sargent DJ, Wieand HS, Fleshman J, Anvari M, et al. A comparison of laparoscopically assisted and open colectomy for colon cancer. *N Engl J Med*. 2004;350(20):2050-9.
18. Berends F, Kazemier G, Bonjer H, Lange J. Subcutaneous metastases after laparoscopic colectomy. *Lancet*. 1994;344(8914):58.
19. Reilly WT, Nelson H, Schroeder G, Wieand HS, Bolton J, O'Connell MJ. Wound recurrence following conventional treatment of colorectal cancer. *Dis Colon Rectum*. 1996;39(2):200-7.
20. Abboudi H, Khan MS, Aboumarzouk O, Guru KA, Challacombe B, Dasgupta P, et al. Current status of validation for robotic surgery simulators—a systematic review. *BJU Int*. 2013;111(2):194-205.

21. Leroy J, Jamali F, Forbes L, Smith M, Rubino F, Mutter D, et al. Laparoscopic total mesorectal excision (TME) for rectal cancer surgery: long-term outcomes. *Surg Endosc.* 2004;18(2):281-9.
22. Pigazzi A, Ellenhorn J, Ballantyne G, Paz I. Robotic-assisted laparoscopic low anterior resection with total mesorectal excision for rectal cancer. *Surg Endosc.* 2006;20(10):1521-5.
23. Delaney CP, Lynch CA, Senagore AJ, Fazio VW. Comparison of robotically performed and traditional laparoscopic colorectal surgery. *Dis Colon Rectum.* 2003;46(12):1633-9.
24. Cahill RA. Natural orifice transluminal endoscopic surgery – here and now. *Surgeon.* 2010;8(1):44-50.
25. Swanström L. Natural orifice transluminal endoscopic surgery. *Endoscopy.* 2009;41(1):82-5.
26. Sodergren MH, Clark J, Athanasiou T, Teare J, Yang G-Z, Darzi A. Natural orifice transluminal endoscopic surgery: critical appraisal of applications in clinical practice. *Surg Endosc.* 2009;23(4):680-7.
27. Wilson J, Dogiparthi K, Hebblethwaite N, Clarke M. Laparoscopic right hemicolectomy with posterior colpotomy for transvaginal specimen retrieval. *Colorectal Dis.* 2007;9(7):662.
28. Raju GS, Pham B, Xiao S-Y, Brining D, Ahmed I. A pilot study of endoscopic closure of colonic perforations with endoclips in a swine model. *Gastrointest Endosc.* 2005;62(5):791-5.
29. Nishimura J, Nakajima K, Omori T, Takahashi T, Nishitani A, Ito T, et al. Surgical strategy for gastric gastrointestinal stromal tumors: laparoscopic vs. open resection. *Surg Endosc.* 2007;21(6):875-8.
30. Rattner D, Kalloo A. ASGE/SAGES working group on natural orifice transluminal endoscopic surgery. *Surg Endosc.* 2006;20(2):329-33.
31. Saad S, Hosogi H. Laparoscopic left colectomy combined with natural orifice access: operative technique and initial results. *Surg Endosc.* 2011;25(8):2742-7.
32. Targarona EM, Gomez C, Rovira R, Pernas JC, Balague C, Guarner-Argente C, et al. NOTES-assisted transvaginal splenectomy: the next step in the minimally invasive approach to the spleen. *Surg Innov.* 2009;16(3):218-22.
33. Baekelandt J, De Mulder PA, Le Roy I, Mathieu C, Laenen A, Enzlin P, et al. Postoperative outcomes and quality of life following hysterectomy by natural orifice transluminal endoscopic surgery (NOTES) compared to laparoscopy in women with a non-prolapsed uterus and benign gynaecological disease: a systematic review and meta-analysis. *Eur J Obstet Gynecol Reprod Biol.* 2017;208:6-15.
34. Marescaux J, Dallemagne B, Perretta S, Wattiez A, Mutter D, Coumaros D. Surgery without scars: report of transluminal cholecystectomy in a human being. *Arch Surg.* 2007;142(9):823-6; discussion 826-7.