

Breast Self-Examination Practice among Female Undergraduate Students in Debre Tabor University, Northcentral Ethiopia: Based on Health Belief Model

Wubet Taklual**, MPH, Aragaw Tesfaw*, MPH, Maru Mekie**, MPH, Tesfaneh Shemelis*, MPH

*Department of Public Health, Debre Tabor University, College of Health Science, Debre Tabor, Ethiopia

**Department of Midwifery, Debre Tabor University, College of Health Science, Debre Tabor, Ethiopia

Please cite this article as: Taklual W, Tesfaw A, Mekie M, Shemelis T. Breast self-examination practice among female undergraduate students in Debre Tabor University, northcentral Ethiopia: Based on health belief model. Middle East J Cancer. 2021;12(4):563-72. doi: 10.30476/mejc.2021.84950.1245.

Abstract

Background: We conducted the present study to assess the practice of breast self-examination (BSE) among Debre Tabor University female undergraduate students, northcentral Ethiopia using health belief model (HBM).

Method: The current institution-based cross-sectional study was carried out among a total of 341 students. Simple random sampling technique was employed to select the study participants. We also used self-administered pretested questionnaires and constructs of HBM for collecting data. Binary and multivariable logistic regression were utilized to identify BSE-associated factors. Statistical significance was stated at P value < 0.05 .

Results: Approximately 45% of the participants had a fair level of knowledge about breast cancer (BC), but only 17% were actually performing BSE practice. The odds of practicing BSE were found to be higher among the participants who had information about BSE practice adjusted odds ratio (AOR)=7.21, 95% confidence interval (CI): (2.46, 21.15)), perceived susceptibility (AOR=14.18, 95% CI:(4.00, 50.48)), self-efficacy (AOR=3.07, 95% CI: (1.09, 8.70)), cue to action (AOR=3.68, 95% CI: (1.17, 11.56)), and net benefit (AOR=7.75, 95% CI: (1.56, 38.55)) compared with counterparts. Whereas, the odds of practicing BSE were found to be lower among those who had poor knowledge of BC (AOR=0.08, 95% CI: (0.03, 0.23)) compared with counterparts.

Conclusion: In this study, knowledge regarding BC and BSE practice was observed to be low. Knowledge about BC, having information on BSE, perceived susceptibility, self-confidence, and cue to action were found to be independent predictors of BSE practice. Providing targeted information about BSE is the best method of changing the behavioral intention of university students about BC and BSE practice.

Keywords: Practice, Breast self-examination, Breast cancer, Health belief model, Ethiopia

*Corresponding Author:

Wubet Taklual, MPH
Department Public Health,
College of Health Sciences,
Debre Tabor University,
Debre Tabor, Ethiopia
Email: wubetaktual21@yahoo.com

Introduction

Cancer, as global public health issue, is one of the major non-communicable diseases and the second leading cause of death in high-income countries and 3rd leading cause of death in low- and middle-income countries.¹ Breast cancer is known to be one of the major malignant tumors that starts in the cells of the breast and affects women mostly.² Worldwide, breast cancer accounts for almost 1 in 4 cancer cases among women and the leading cause of cancer death in over 100 countries. It ranks third in terms of incidence, but second in terms of mortality.³ It is the most frequently diagnosed malignancy among women next to lung cancer. It alone accounts for 25% of all cancer cases and 15% of all cancer deaths among females.⁴

Even though a higher prevalence is observed in developed countries, the mortality due to breast cancer is higher in developing ones. Almost 50% of breast cancer cases and 58% of deaths occur in developing countries (LMICs).⁵ Studies have also shown that the highest age-standardized breast cancer mortality rate is recorded in Africa, specifically in Sub-Saharan Africa.^{4, 6}

Like other Sub-Saharan African countries, the numbers of new breast cancer cases are increasing alarmingly in Ethiopia. It has an estimated age standardized incidence rate of 43 cases per 100,000 women and has become the most prevalent cancer in this country with an increasing trend and advanced stage presentations.^{7, 8} According to the Addis Ababa city cancer registry report, breast cancer accounts for 33% of all cancer cases in women.⁹

About 30% - 50% of cancers could be prevented by avoiding risk factors and implementing evidence-based prevention strategies through early detection of cancer and management of patients who develop cancer.¹ Similarly, early detection of breast cancer plays a key role in the survival of breast cancer patients.¹⁰ According to evidence, patients diagnosed at early stages have demonstrated better survival than those diagnosed at late stages.¹¹ However, breast cancer awareness and knowledge of the benefits associated with early detection

and diagnosis are poor in most Sub-Saharan African countries and consequently, advanced stage presentation remains a common feature in the region.¹²⁻¹⁴

Several studies have reported that knowledge and practice of breast self examination (BSE) and clinical breast examination increase the likelihood of presenting at early stages with a high chance of being cured.^{15, 16} However, studies in Africa have indicated that the practice of BSE is low among university students. In Cameroon, only 3% performed BSE and 9% know how to perform BSE. Similarly, 19% of studied cases performed BSE in Nigeria. Lack of knowledge about BSE is mentioned as a significant barrier to practicing it.^{17, 18} Despite sufficient knowledge about the importance of BSE, only 23% of medical students performed BSE in Haramaya University. BSE is the best tool to diagnose breast cancer early in low-income countries like Ethiopia where access to mammography is limited.¹⁹

In Ethiopia, the attention of the government and health care systems were towards communicable diseases rather than cancer in previous years, but currently, in Ethiopia, there is a plan to control the increasing patterns of breast cancer through screening and early detection programs, such as mammography, clinical, and BSE.²⁰ However, there is the lack of evidence concerning the level of BSE practice among university students in this country. Studies in this regard are generally very scarce. In addition, predicting student's behavioral intention for performing BSE is of great importance for a country like Ethiopia, where there is poor health care delivery system. Thus, understanding students' beliefs on BSE practice is crucial for evidence-based behavioral intervention among university students. Therefore, we aimed to assess the practice of BSE among Debre Tabor University female students using health belief model (HBM).

Methods

Study setting and period

The present institution-based cross-sectional study was carried out at Debre Tabor University

(DTU) in December 2018. This university is one of the higher education institutions in Ethiopia located in the northcentral part of the country, 667 km from the capital. It is established in 2008 in Debre Tabor town, South Gondar administrative zone of Amhara regional state at the foot of Guna Mountain. The University has unique and different features by implementing integrative Hybrid Innovative Curriculum, particularly in medical and health sciences disciplines.²¹ There are five faculties and one college in the university. At the time of the study, there were more than 12,000 students following their education in a regular program.

Study population and inclusion criteria

Our study population comprised all the regular undergraduate female students. Female students below the age of 18, health science students, and students who lived outside the university were excluded. Health science students were excluded from the study since they have some basic information about breast cancer and BSE in their courses.

Sample size and sampling procedure

The sample size of the study was determined considering the following assumptions: taking a prevalence of BSE practice from a study conducted at Debre Berhan University 28.3%,²² a 5% margin of error, a confidence level (CI) of 95% and considering non-response rate of 10%, the representative sample of participants was determined (n=342) using a single population proportion formula for a cross-sectional survey in Epi info version 7.2. Therefore, our final sample size was 342 students. A simple random sampling technique was utilized to select the study participants. After determining proportional allocation of the sample to the population size, the sampling frame of female students in each faculty was prepared and they were selected from the frame using simple random sampling method.

Data collection procedures and quality assurance

The data was collected with structured self-administered questionnaire developed by reviewing different literatures.^{17, 22, 23} The questionnaire was first prepared in English, then translated to Amharic, and finally translated back into English.

Table 1. Socio-demographic characteristics of female students of Debre Tabor University, northcentral Ethiopia

Variable	Frequency	Percent
Age in years		
18-19	27	7.9
20-24	305	89.2
25-29	10	2.9
Religion		
Orthodox	280	81.9
Muslim	35	10.2
Protestant	27	7.9
Marital status		
Single	279	81.6
Married	50	14.6
Separated	13	3.8
Department		
Social Sciences	151	44.2
Natural Sciences	191	55.8
Year of study		
Second	136	39.8
Third and above	206	60.2
Information about breast cancer		
Yes	192	56.1
No	150	43.9
Source of information		
Mass media	93	48.4
Books/magazine	51	26.6
Health facility	47	24.5
Health professional	73	38.0
Family/friends	77	40.1

The components of the questionnaire were socio-demographic characteristics, perceived severity, susceptibility, benefits, cues to action, barriers, and self-efficacy for performing BSE. The questionnaire was given to the participants so that fill it in a private setting after taking their consent. Four health officers and three masters of public health professionals participated in the facilitation and supervision of the data collection process, respectively.

To maintain the quality of the data, the questionnaires were prepared in a simple and easily understandable language. A two-day training was performed for the facilitators on the data collection process. The questionnaires were pretested in 5% of the sample population and the necessary modifications were done. The pretested data were not included in the final analysis. Strict supervision and monitoring was done during data collection.

Table 2. Responses of Debre Tabor University female students to HBM scale, northcentral Ethiopia, 2019

Scales	Agree		Neutral		Disagree	
	Frequency	Percent	Frequency	Percent	Frequency	Percent
Susceptibility	190	55.6	37	10.8	115	33.6
Seriousness	109	31.9	17	5.0	216	62.3
Benefit of BSE	94	27.5	37	10.8	160	46.8
Barriers of BSE	188	55.0	18	5.3	136	39.8
Self-efficacy	154	45.0	20	5.8	168	49.1
Health motivation	Yes N (%)				No N (%)	
	239 (69.9)				103 (30.1)	

BSE: Breast self-examination; HBM: Health belief model

Measurements

For the constructs of HBM, five item responses were prepared for each construct (strongly agree score's 5 points) to strongly disagree (scores "1" point). Perceived susceptibility of breast cancer consisted of three items scored from 3 to 15. Seriousness of breast cancer comprised six items scored from 6 to 30. Benefits consisted of four items scored from 4 to 20. BSE barriers included eight items scored from 8 to 40. BSE self-efficacy consisted of nine items scored from 9 to 45, and cues to action comprised five items with 'Yes or No' questions. For all the constructs of HBM, higher scores indicated having high perception towards performing BSE; whereas, higher score of barriers indicated high barrier to perform BSE. Regarding perceived net benefit, we employed the sum score of perceived benefit minus that of perceived barriers.²⁴ The reliability coefficient for each HBM constructs was calculated using Cronbach's alpha. The reliability coefficient was 0.813, 0.831, 0.770, 0.850, and 0.860 for susceptibility, seriousness, benefit, barrier, and self-efficacy, respectively.

BSE practice was assessed via 'Yes' or 'No' type questions. The subjects who responded "Yes" for a question like "have you ever performed BSE for screening cancer?" were considered as those practicing BSE.

Data Processing and Analysis Procedures

The data were entered in EpiData version 3.1 and cleaned and analyzed with SPSS version 23. Descriptive statistics were computed using frequency and other statistical summary measures. A chi-square test on top of bivariable and multivariable logistic regression analysis were

used to identify the factors associated with BSE practice. The crude and adjusted odds ratios (OR) were used to identify the significance, strength, and direction of association at 95% CI. A *P*-value of <0.05 was employed to decide whether the association was significant.

Ethical considerations

An ethical approval was obtained from the Research Ethics Committee of Debre Tabor University under a reference number of R.No: CHS/5173/2019. Oral informed consent was obtained from the students prior to the study, after a brief explanation about the purpose of the study. Participation in the study was fully voluntary and all the information collected from the participants were kept confidential.

Result

Socio-demographic characteristics of the study participants

The mean age of the subjects was 21.18 ± standard deviation (SD) 1.51 years. More than 80% of the study participants were orthodox Christian religion followers. With regards to department distribution, more than half, 191 (55.8%) of the study participants, were from natural sciences and the rest, 151 (44.2%), were from social sciences stream.

More than half of the study participants, 192 (56.1%), were familiar with breast cancer. The main source of information about breast cancer in 93 participants (48.4 %) was mass media followed by family/friends in 77 (40.1%) of them, and health professionals in 73 (38.0%) (Table 1).

Table 3. Differences in health-associated beliefs with regards to practice of BSE among female Debre Tabor University, northcentral Ethiopia, 2019

Scales	No BSE practice	BSE practice	X ² -value	P value
Susceptibility				
Agree	144 (50.7)	46 (79.3)	17.06	0.000
Unsure	32 (11.3)	5 (8.6)		
Disagree	108 (38.0)	7 (12.1)		
Seriousness				
Agree	78 (27.5)	31 (53.4)	15.0	0.001
Unsure	15 (5.3)	2 (3.4)		
Disagree	191 (67.3)	25 (43.1)		
Benefit				
Agree	67 (28.2)	27 (50.9)	11.09	0.004
Unsure	34 (14.3)	3 (5.7)		
Disagree	137 (57.6)	23 (43.4)		
Barrier				
Agree	172 (60.6)	16 (27.6)	22.79	0.000
Unsure	15 (5.3)	3 (5.3)		
Disagree	97 (34.2)	39 (67.2)		
Self-efficacy				
Agree	115 (40.5)	39 (67.2)	14.30	0.001
Unsure	17 (6.0)	3 (5.2)		
Disagree	152 (53.5)	16 (27.6)		
Cue to action				
Yes	191 (67.3)	48 (82.8)	5.50	0.019
No	93 (32.7)	10 (17.2)	0.019	
Net benefit				
Positive	9 (3.2)	19 (32.8)	56.10	0.000
Negative	275 (96.8)	39 (67.2)	0.000	

BSE: Breast self-examination

HBM

Table 2 represents the summary of health belief scale categorized into three sub scales. More than half 190 (55.6%) of the participants believed that they are susceptible to breast cancer. Meanwhile, 115 (33.6%) of them believed that they are not susceptible to breast cancer. More than half, 216 (62.3%) of the participants, believed that breast cancer is not serious and 17 (5.0%) were unsure about the seriousness of breast cancer. Nearly half of the study participants, 160 (46.7%), did not believe that BSE is beneficial and 94 (27.5%) acknowledged the advantage of BSE. 154 (45.0%) of the participants had the confidence to perform BSE. More than half of the participants, 188 (55.0%), believed that barriers prevented them from performing BSE. 70% of the study participants had the motive to perform BSE (Table 2).

Differences in health beliefs among study participants

More than three quarters (79.9%) of the study participants who practices BSE agreed that they are susceptible to breast cancer. On the other hand, only half of the non-practicing subjects agreed on their susceptibility to breast cancer. 172 (60.6%) of the practicing and 16 (27.6%) of the non-practicing ones believed that barriers prevented them from practicing BSE (Table 3).

Factors associated with practice of breast self-examination

Information about breast cancer, knowledge, perceived susceptibility, self-efficacy, cues to action, and net benefit were found to be statistically significant with the practice of BSE in the final model. The multivariable analysis revealed that the odd of practicing BSE was found to be seven times higher among the participants who had information about BSE (adjusted odds

Table 4. Multivariable analysis of the factors affecting BSE practice among female students in Debre Tabor University, northcentral Ethiopia, 2019

Variables	Practice of BSE		COR (95% CI)	AOR (95% CI)
	No N (%)	Yes N (%)		
Department				
Social	117 (41.2)	34 (58.6)	2.02 (1.14, 3.59)	1.88 (0.69, 5.15)
Natural	167 (58.8)	24 (41.4)	1	1
Information on BSE				
Yes	77 (27.1)	33 (56.9)	3.55(1.98, 6.35)	7.21 (2.46, 21.15)*
No	207 (72.9)	25 (43.1)	1	1
Knowledge on BCA				
Poor	181 (63.7)	7 (12.1)	0.08 (0.03, 0.18)	0.08 (0.03, 0.23)*
Good	103 (36.6)	51 (87.9)	1	1
Susceptibility				
Agree	144 (50.7)	46 (79.3)	4.93 (2.14, 11.34)	14.18 (4.00, 50.48)*
Unsure	32 (11.3)	5 (8.6)	2.41(0.72, 8.11)	1.55 (0.22, 10.93)
Disagree	108 (38.0)	7 (12.1)	1	1
Seriousness				
Agree	78 (27.5)	31 (53.4)	3.04(1.69, 5.47)	2.51 (0.97, 6.52)
Unsure	15 (5.3)	2 (3.4)	1.02 (0.22, 4.72)	0.34 (0.02, 6.10)
Disagree	191 (67.3)	25 (43.1)	1	1
Benefit				
Agree	67 (28.2)	27 (50.9)	2.40 (1.28, 4.50)	0.81 (0.27, 2.40)
Unsure	34 (14.3)	3 (5.7)	0.53 (0.15, 1.85)	0.17 (0.02, 1.50)
Disagree	137 (57.6)	23 (43.4)	1	1
Barrier				
Agree	172 (60.6)	16 (27.6)	0.23 (0.12, 0.44)	0.49 (0.17, 1.39)
Unsure	15 (5.3)	3 (5.3)	0.50 (0.14, 1.81)	2.09 (0.25, 17.19)
Disagree	97 (34.2)	39 (67.2)	1	1
Self-efficacy				
Agree	115 (40.5)	39 (67.2)	3.22 (1.72, 6.05)	3.07 (1.09, 8.70)*
Unsure	17 (6.0)	3 (5.2)	1.68 (0.44, 6.35)	2.16 (0.37, 12.57)
Disagree	152 (53.5)	16 (27.6)	1	1
Cue to action				
Yes	191 (67.3)	48 (82.8)	2.34 (1.13, 4.83)	3.68 (1.17, 11.56)*
No	93 (32.7)	10 (17.2)	1	1
Net benefit				
Positive	9 (3.2)	19 (32.8)	14.89 (6.29, 35.22)	7.75 (1.56, 38.55)*
Negative	275 (96.8)	39 (67.2)	1	1

AOR: Adjusted Odds Ratio; BCA: Breast Cancer; BSE: Breast self-examination; COR: Crude Odds Ratio; CI: Confidence Interval; * significant in the final model

ratio ((AOR) = 7.21, 95% CI: (2.46, 21.15)). There was a significant association between breast cancer knowledge and BSE. The odds of practicing BSE was found to be low among those who had poor knowledge (AOR= 0.08, 95% CI: (0.03, 0.23)) compared with counterparts. Meanwhile, the odds of practicing BSE was found to be higher among the participants with perceived susceptibility (AOR=14.18, 95% CI: (4.00, 50.48)), self-efficacy (AOR=3.07, 95% CI: (1.09, 8.70)), cues to action (AOR= 3.68, 95%, CI: (1.17, 11.56)), and net benefit of BSE (AOR=7.75,

(1.56, 38.55)) (Table 4).

Discussion

In this study, 17% of the students were found to practice BSE, despite 40% of those who had information about it. The finding of the present study is consistent with the previous researches,²⁵⁻²⁷ but the proportion of BSE practice among students was lower than the studies conducted among female health science students in Ethiopia (23%), Nigeria (25%), and Uganda (43.6%).^{22, 23, 28, 29} The difference could be ascribed to the

level of awareness about breast cancer and screening. Moreover, the differences might be due to health sciences students who could have more information about breast cancer were excluded in this study unlike previous studies.

In this work, the participants with information about BSE were 7.21 times more likely to practice BSE compared with counterparts (AOR=7.21, 95% CI; (2.46, 21.5)). This implied that information about BSE is a precursor for practice. Therefore, IEC through different channels is very important for increasing the screening practice of breast cancer. Similarly, those with poor knowledge on breast cancer were less likely to practice BSE than the counterparts (AOR=0.08, 95 % CI;(0.03, 0.23)). This finding is consistent with that of previous studies.²⁹⁻³¹ This is explained through the fact that knowledge on breast cancer is precondition for screening and early diagnosis. This is supported by studies conducted in different countries.³²⁻³⁴ In our study, only a third of the participants were knowledgeable about breast cancer risk factors and clinical picture. This shows that the students have poor knowledge about BC as well as screening practice. Therefore, the government should take some measures regarding university students since they are a better medium for IEC, while they are back to the community during vocation or after graduation when they are employed at different institutions. Behavioral change in communication among university students through continuous training, peer education, and health education in universities are pivotal strategies for early detection and management of breast cancer. Furthermore, it is recommended to incorporate reproductive health (family health) course to other non-health faculty students.

In the present research, in the HBM constructs, perceived susceptibility, self-efficacy, and cue to action were the predictors of BSE practice; whereas, perceived benefit, barrier, and seriousness were not found to be significantly associated with BSE practice. In this study, the participants who confirmed their susceptibility to breast cancer were 14.18 times more likely to practice BSE than the counterparts (AOR=14.18, 95% CI (4.00,

50.48)). Our finding also revealed that about 80% of the BSE performers agreed on susceptibility to breast cancer; whereas, among BSE non-performers, only 50% supported future breast cancer susceptibility. This could explain that those who perceived they are susceptible believe that BSE would help to detect breast cancer early and improve the outcome. This finding is in accordance with that of other studies.^{24, 27, 35}

The bi-variable analysis revealed that the odds of practicing BSE was three times higher among the participants who agreed on perceived severity (COR=3.04, 95% CI:(1.69, 5.47)). However the association disappeared when adjusted for confounders. This implied that the perceived degree of threats on breast cancer would improve an individual screening practice. Therefore, giving health education about threats of breast cancer for university students as well as the community at large could improve the beliefs on screening practice, which is a cost-effective strategy for reducing breast cancer incidence and prevalence in low-income countries. This result is supported by a study conducted in Ethiopia among female teachers.²⁴

Perceived benefit also illustrated to have a significant association with BSE practice based on bivariate analysis. Those participants who agreed on perceived benefit of BSE were 2.4 times more likely to practice BSE compared with their counterparts (COR=2.4, 95% CI: (1.28, 4.50)) though the association vanished when adjusted for confounders. Those with better understanding about perceived benefit of screening were more likely to practice BSE. This finding is consistent with that of previous studies.^{24, 26, 36, 37}

Herein, self-efficacy was also significantly associated with BSE practice. The odds of performing BSE was found to be three times higher among self-confident participants (AOR= 3.07, 95% CI :(1.09, 8.70)). This finding is consistent with previously indicated results.^{27, 30, 36, 38}

This indicated that one's confidence to successfully perform an action significantly affect their behavioral practice. Therefore, providing

information or training to the students would affect the practice. Similarly, in the present study those participants with a motivation towards performing BSE were 3.68 times more likely to practice BSE than the counterparts (AOR=3.68, 95% CI:(1.17, 11.56)), which is consistent with previous studies.^{35, 38} This implies that motivating an individual will have an intention to perform particular behavior. Therefore, cues are a good strategy to activate or stimulate BSE practice. Thus, a reminder of BSE using different methods would improve their practice.

The net benefit was also significantly associated with practicing BSE (AOR=7.75, 95% CI: (1.56, 38.55)). This implied that the students have an intention to develop BSE behavior in the future. Therefore, giving information about breast cancer and BSE will improve screening practice and early diagnosis of breast cancer. This finding was in agreement with the constructs of HBM in which those participants with a high score of perceived susceptibility, perceived severity, perceived benefit, self-efficacy, and cue to action and a low score of perceived barriers were more likely to practice BSE.^{39, 40}

The study had certain limitations. Initially, it included only female educated university students; this makes it very difficult to generalize the results to a large group of population, where literacy rate and rural community are very low. Secondly, since the data was collected via self-administer questionnaire, the respondents may have recalled bias on BSE performance (Since BSE practice was assessed by the pervious performance of the respondents, the may not remember their pervious performance), unconscious response, and dishonest answer. In addition, the HBM constructs is rather descriptive than explanatory; accordingly, it is better to integrate with other models which contains environmental context and strategies for changing human behavior.

Conclusion

In conclusion, the practice of BSE was low. The knowledge on breast cancer, having information on BSE, perceived susceptibility,

self-confidence, and cue to action were found to be the independent predictors of BSE practice. Providing information to the students through different channels is very important to improve the knowledge in this regard and practice of BSE. It is highly recommended that the government of Ethiopia consider incorporating reproductive and family health courses to other non-health faculty students.

Availability of Data and Materials

The datasets utilized in this work were provided by the corresponding author and could be accessible through reasonable request.

Acknowledgment

The authors would like to acknowledge the College of Health Science, Debre Tabor University for securing the ethical review process. The authors should like to express the deepest gratitude to the supervisors, data collectors, and study participants.

Conflict of Interest

None declared.

References

1. WHO. Cancer fact sheet, 2017.[Internet] Available at: <http://www.who.int/en/>. (Accessed date: December, 2018).
2. American Cancer Society. Breast Cancer Facts & Figures 2015-2016. Atlanta: American Cancer Society, Inc. 2015. [Internet] Available at: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/bre> .(Accessed date:December,2018).
3. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68(6):394-424. doi:10.3322/caac.21492.
4. Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, et al. Cancer incidence and mortality worldwide: Sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2015; 136(5):E359-86.doi: 10.1002/ijc.29210.
5. World Breast Cancer Report. International Prevention Research Institute, 2012. [Internet] Available at: <https://i-pri.org/2012/11/04/world-breast-cancer-report-2012/>. (Accessed date, December, 2018)

6. Azubuike SO, Muirhead C, Hayes L, McNally R. Rising global burden of breast cancer: the case of sub-Saharan Africa (with emphasis on Nigeria) and implications for regional development: a review. *World J Surg Oncol*. 2018;16(1):63. doi:10.1186/s12957-018-1345-2.
7. Tessema S, Kifle M, Asefa M, Abayneh G, Tsegaye B, Woldetinsae M, et al. Estimates of cancer incidence in Ethiopia in 2015 using population-based registry data. *J Glob Oncol*. 2018;4:1-11. doi:10.1200/JGO.17.00175.
8. Abate SM, Yilma Z, Assefa M, Tigeneh W. Trends of breast cancer in Ethiopia. *Int J Cancer Res Mol Mech*. 2016;2(1). doi:10.16966/2381-3318.121.
9. American Cancer Society. Breast Cancer Facts & Figures 2017-2018. Atlanta: American Cancer Society, Inc. 2017. [Internet] Available at: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/breast-cancer-facts-and-figures/breast-cancer-facts-and-figures-2017-2018.pdf>. (Accessed date, November, 2018).
10. Yip CH, Smith RA, Anderson BO, Miller AB, Thomas DB, Ang ES, et al. Guideline implementation for breast healthcare in low- and middle-income countries: early detection resource allocation. *Cancer*. 2008;113(8 Suppl):2244-56. doi:10.1002/cncr.23842.
11. Kantelhardt EJ, Zerche P, Mathewos A, Trocchi P, Addissie A, Aynalem A, et al. Breast cancer survival in Ethiopia: A cohort study of 1,070 women. *Int J Cancer*. 2014;135(3):702-9. doi:10.1002/ijc.28691.
12. Jedy-Agba E, McCormack V, Adebamowo C, Dos-Santos-Silva I. Stage at diagnosis of breast cancer in sub-Saharan Africa: a systematic review and meta-analysis. *Lancet Glob Health*. 2016;4(12):e923-e935. doi:10.1016/S2214-109X(16)30259-5.
13. Pruitt L, Mumuni T, Raikhel E, Ademola A, Ogundiran T, Adenipekun A, et al. Social barriers to diagnosis and treatment of breast cancer in patients presenting at a teaching hospital in Ibadan, Nigeria. *Glob Public Health*. 2015;10(3):331-44. doi:10.1080/17441692.2014.974649.
14. Dickens C, Joffe M, Jacobson J, Venter F, Schüz J, Cubasch H, et al. Stage at breast cancer diagnosis and distance from diagnostic hospital in a periurban setting: a South African public hospital case series of over 1,000 women. *Int J Cancer*. 2014;135(9):2173-82. doi:10.1002/ijc.28861.
15. Stapleton MJ, Mullan BP, Dey S, Hablas A, Gaafar R, Seifeldin IA, et al. Patient-mediated factors predicting early and late stage presentation of breast cancer in Egypt. *Psychooncology*. 2011;20(5):532-7. doi: 10.1002/pon.1767.
16. Romanoff A, Constant HT, Johnson MK, Guadamos MC, Vega AMB, Zunt J, et al. Association of previous clinical breast examination with reduced delays and earlier-stage breast cancer diagnosis among women in Peru. *JAMA Oncol*. 2017. doi: 10.1001/jamaoncol.2017.1023.
17. Nde FP, Assob JC, Kwenti TE, Njunda AL, Tainenbe TR. Knowledge, attitude and practice of breast self-examination among female undergraduate students in the University of Buea. *BMC Res Notes*. 2015;8:43. doi: 10.1186/s13104-015-1004-4.
18. Gwarzo UM, Sabitu K, Idris SH. Knowledge and practice of breast-self examination among female undergraduate students of Ahmadu Bello University Zaria, northwestern Nigeria. *Ann Afr Med*. 2009;8(1): 55-8. doi:10.4103/1596-3519.55766.
19. Anderson OB, Cazap E, El Saghir SN, Yip C, Khaled HM, Otero IV, et al. Optimisation of breast cancer management in low-resource and middle-resource countries: executive summary of the Breast Health Global Initiative consensus, 2010. *Lancet Oncol*. 2011; 12:387-98. doi: 10.1016/S1470-2045(11)70031-6.
20. FMOH. National Cancer Control Plan (2016-2020). Disease prevention and control directorate. Addis Ababa, Ethiopia: Federal Ministry of Health ; 2015. [Internet] Available at: [https://www.iccpportal.org/sites/default/files/plans/NC CP%20Ethiopia%20Final%2020261015.pdf](https://www.iccpportal.org/sites/default/files/plans/NC%20CP%20Ethiopia%20Final%2020261015.pdf). (Accessed date: November, 2018).
21. Asefa M. Introducing innovative curriculum. Addis Ababa, Ethiopia Debre Tabor University 2012. [Internet] Available at: <http://www.dtu.edu.et/academic-quality-assurance/> (Accessed date: November, 2018).
22. Birhane K, Alemayehu M, Anawte B, Gebremariyam G, Daniel R, Addis S, et al. Practices of breast self-examination and associated factors among female Debre Berhan University students. *Int J Breast Cancer*. 2017;2017:8026297. doi:10.1155/2017/8026297.
23. Ameer K, Abdulie MS, Pal KS, Arebo K, Kassa GG. Breast cancer awareness and practice of breast self-examination among female medical students in Haramaya University, Harar, Ethiopia. *IJIMS*. 2014;2(2):109-19.
24. Birhane N, Mamo A, Girma E, Asfaw S. Predictors of breast self - examination among female teachers in Ethiopia using health belief model. *Arch Public Health*. 2015;73(1):39. doi: 10.1186/s13690-015-0087-7.
25. Nemenqani DM, Abdelmaqsood SH, Al-Malki A-HA, Orajja AA, Al-Otaibi EM. Knowledge, attitude and practice of breast self examination and breast cancer among female medical students in Taif, Saudi Arabia. *Open J Prev Med*. 2014;4(2):69-77.
26. Gençtürk N, Demirezen E, Ay F. Health beliefs of midwifery students at Istanbul University about breast cancer and breast self-examination acknowledgements. *J Cancer Educ*. 2017;32(4):784-9. doi: 10.1007/s13187-016-1014-5.
27. Akhtari-Zavare M, Juni MH, Ismail IZ, Said SM, Latiff LA. Health beliefs and breast self-examination among undergraduate female students in public

- universities in Klang Valley, Malaysia. *Asian Pac J Cancer Prev*. 2015;16(9):4019-23. doi: 10.7314/apjcp.2015.16.9.4019.
28. Godfrey K, Agatha T, Nankumbi J. Breast cancer knowledge and breast self-examination practices among female university students in Kampala, Uganda: A descriptive study. *Oman Med J*. 2016;31(2):129-34. doi: 10.5001/omj.2016.25.
 29. Alwan N, Al-Diwan J, Al-Attar W, Eliessa R. Knowledge, attitude & practice towards breast cancer & breast self-examination in Kirkuk University, Iraq. *Asian Pacific Journal of Reproduction*. 2012;1(4):308-11. doi: 10.1016/S2305-0500(13)60098-6.
 30. Abolfotouh MA, BaniMustafa AA, Mahfouz AA, Al-Assiri MH, Al-Juhani AF, Alaskar AS. Using the health belief model to predict breast self examination among Saudi women. *BMC Public Health*. 2015;15:1163. doi: 10.1186/s12889-015-2510-y.
 31. Dündar PE, Ozmen D, Oztürk B, Haspolat G, Akyildiz F, Coban S, et al. The knowledge and attitudes of breast self-examination and mammography in a group of women in a rural area in western Turkey. *BMC Cancer*. 2006;6:43. doi: 10.1186/1471-2407-6-43.
 32. Joffe M, Ayeni O, Norris SA, McCormack VA, Ruff P, Das I, et al. Barriers to early presentation of breast cancer among women in Soweto, South Africa. *PLoS One*. 2018;13(2):e0192071. doi: 10.1371/journal.pone.0192071.
 33. Akuoko CP, Armah E, Sarpong T, Quansah DY, Amankwaa I, Boateng D. Barriers to early presentation and diagnosis of breast cancer among African women living in sub-Saharan Africa. *PLoS One*. 2017. 12(2):e0171024. doi:10.1371/journal.pone.0171024.
 34. Parsa P, Kandiah M, Abdul Rahman H, Zulkefli NM. Barriers for breast cancer screening among Asian women: a mini literature review. *Asian Pac J Cancer Prev*. 2006;7(4):509-14.
 35. Canbulat N, Uzun O. Health beliefs and breast cancer screening behaviors among female health workers in Turkey. *Eur J Oncol Nurs*. 2008;12(2):148-56. doi: 10.1016/j.ejon.2007.12.002.
 36. Noroozi A, Jomand T, Tahmasebi R. Determinants of breast self-examination performance among Iranian women: an application of the health belief model. *J Cancer Educ*. 2011;26(2):365-74. doi: 10.1007/s13187-010-0158-y.
 37. Asghari E, Nahamin M, Khoshtarash M, Ghanbari A, Parizad N, Mahdavi N, et al. The relationship between health belief and breast self-examination among Iranian university students. *International Journal of Women's Health and Reproduction Science (IJWHR)*. 2016;4(3): 110-23.
 38. Avci IA. Factors associated with breast self-examination practices and beliefs in female workers at a Muslim community. *Eur J Oncol Nurs*. 2008;12(2):127-33. doi: 10.1016/j.ejon.2007.11.006.
 39. Champion VL. Use of the health belief model in determining frequency of breast self-examination. *Res Nurs Health*. 1985;8(4):373-9. doi: 10.1002/nur.4770080410.
 40. Champion VL. The relationship of breast self-examination to health belief model variables. *Res Nurs Health*. 1987;10(6):375-82. doi: 10.1002/nur.4770100605.